

STA 321

Spring 2016

Lecture 12

Tuesday, March 1

- **Hypothesis Tests (Ch 9.1)**
- **Two types of errors (Ch 9.2)**

Hypothesis Testing

- Fact: It is easier to *prove* that a parameter **isn't** equal to a particular value than it is to prove it **is** equal to a particular value
- Hypothesis testing: *Proof by contradiction*:
 - we set up the belief we wish to disprove as the **null hypothesis** (H_0) and the belief we wish to prove as our **alternative hypothesis** (H_1) (or: research hypothesis)

Analogy: Court trial

- In American court trials, the jury is instructed to think of the defendant as innocent:

H_0 : Defendant is innocent

- District attorney, police involved, plaintiff, etc., bring every evidence, hoping to prove

H_1 : Defendant is guilty

- Which hypothesis is correct?
- Does the jury make the right decision?

Back to statistics ...

Critical Concepts

- Two hypotheses: the null and the alternative
- Process begins with the assumption that the null is *true*
- We calculate a test statistic to determine if there is enough evidence to infer that the alternative is true
- Two possible decisions:
 - Conclude there is enough evidence to reject the null (and therefore accept the alternative)
 - Conclude that there is not enough evidence to reject the null
- Two possible errors?

What about those errors?

Two possible errors:

- Type I error: Rejecting the null when we shouldn't have [$P(\text{Type I error}) = \alpha$]
- Type II error: Not rejecting the null when we should have [$P(\text{Type II error}) = \beta$]

Terminology

- Choosing the group corresponding to H_1 is called “rejecting the null hypothesis”
- Choosing the group corresponding to H_0 is called “not rejecting the null hypothesis”

More Terminology

- Note: We set the probability that the correct decision is made when H_0 is true. This probability is typically termed $1-\alpha$.
- Thus, α is the probability of making a mistake when H_0 is true (rejecting when you should NOT reject).
- Once α is set, the cutoff is determined. The most common value for α is 0.05.
- Aside from making α small (it is the chance of a mistake) there is no absolute justification for this choice compared to others.

More Terminology

- The choice of α determines the cutoff, and thus the probability of making the correct decision when H_1 is true (when H_1 is true the correct decision is to reject the null hypothesis).
- This probability is called the **power** of the test.
- Thus, we want $1-\alpha$ and the power to be high (close to 1).

Four Possibilities

- There are two possible states of the world – either the null hypothesis is true or the alternative hypothesis is true.
- You can make two decisions – either reject the null hypothesis or do not reject the null hypothesis.
- Thus, there are four possibilities. Two correspond to correct decisions and two are errors.

The four possibilities, with terminology

	Choose H_0 (“ <u><i>do not reject</i></u> <u><i>H₀</i></u> ”)	Choose H_1 (“ <u><i>reject H₀</i></u> ”)
H_0 is true	Correct answer	WRONG! (<u>Type I error</u>)
H_1 is true	WRONG! (<u>Type II Error</u>)	Correct answer


Type I and Type II errors

- Type I error corresponds to rejecting H_0 when H_0 is in fact true. The probability of making this mistake when H_0 is true is α
- Type II error corresponds to *not* rejecting H_0 when H_1 is in fact true. The probability of making this mistake when H_1 is true is 1 minus the power of the test.

By the way..regression is also useful for this

- Which of the Shakespeare Sonnets were really within the Shakespeare canon?

<http://www.sonnets.org/canon.htm>



Example

- Suppose you have a document that may or may not have been written by author Bill
- You observe that Bill uses a particular form X of the word 80% of the time
- The document in question has 84 instances of the word choice, and word X is used 58 times.

Example continued

- Null hypothesis $H_0: p=0.8$ vs. $H_1 : p \neq 0.8$
- The ***null distribution*** (when H_0 is true) is normal with mean 0.8 and standard deviation $\sqrt{0.8 \cdot 0.2 / 84} = 0.0436$.
- The null distribution is the sampling distribution of the sample statistic when the null hypothesis is assumed true.

Example continued

- We have the null distribution $N(0.8, 0.0436)$
- Let's choose $\alpha=0.05$
- We will reject H_0 for \hat{p} 's far away from 0.8.
- The cutoff to be the 2.5th and the 97.5th percentile of the null distribution. This corresponds to $Z=(-1.96)$ and $Z=1.96$, and a cutoff of
- $(-1.96)(0.0436)+0.8=0.7145$ and
- $(1.96)(0.0436)+0.8=0.8855$

Example continued

- $(-1.96)(0.0436)+0.8=0.7145$ and
- $(1.96)(0.0436)+0.8=0.8855$
- Our ***rejection region*** consists of those values of \hat{p} that are not between 0.7145 and 0.8855
- Our actual observed \hat{p} was $58/84=0.6904$.
- Thus our conclusion is to reject H_0 . We would conclude that H_1 is true and that Bill did not write the document.

Example continued

- How likely is $58/84=0.6904$ under the null hypothesis?
- The z-score is $(0.6904-0.8)/0.0436= -2.51$.
- Beyond (Here="below") a z-score of -2.51 is probability 0.00597.
- All values ***at least as extreme as*** a z-score of -2.51 have together probability $2 \times 0.00597 = 0.0119$.
- This is the ***P-value: The probability, assuming that the null hypothesis is true, of observing anything at least as extreme as what we actually observed.***
- ***"As extreme" = "providing as much, or more evidence against the null hypothesis"***

Example continued

- By construction, the probability of type I error is $\alpha=0.05=5\%$. What is the probability of type II error?
- Type II error is not rejecting H_0 when in fact H_1 is true.
- Assume that another author, Maria, could have written the manuscript, and she chooses form X 60% of the time.
- The sampling distribution for her would be the ***alternative distribution*** (when H_1 is true), normal with mean 0.6 and standard deviation $\text{sqrt}(0.6*0.4/84)=0.0535$.
- We need to find the probability that the alternative distribution places above the cutoff 0.7145.
- The Z-score for a $N(0.6,0.0535)$ is $Z=(0.7145-0.6)/0.0535=2.14$ and the probability of type II error is $0.01609=1.6\%$

Another Example

- A person claims to have ESP.
- You test him/her by having one of four images displayed on a computer screen to a second individual.
- The individual claiming to have ESP has to guess what is on the computer screen.
- Just by random guessing, you would expect the “ESP person” to get 25% of the images correct.
- Thus, a claim of ESP would be strengthened if the person guessed more than 25% of the images correctly.

ESP Example in Statistical Terms

- Suppose you show the person $n=100$ images. Each guess is correct or incorrect (binary, dichotomous).
- The proportion of correct guesses is \hat{p} .
- The true proportion of correct guesses (unknown to us) is p .
- Under random guessing, $p=0.25$
- If the person has ESP, $p>0.25$


Null and Alternative Hypotheses

- The cutoff is based on the null hypothesis.
- $H_0 : p=0.25$ allows us to find a sampling distribution.
- The hypothesis “ $p>0.25$ ” would not determine a “ p ” and a sampling distribution.
- It is a ***composite, one-sided alternative***.
- The alternative hypothesis is the “interesting conclusion” which would get the paper published.

Null Distribution

- Under the null hypothesis, $p=0.25$.
- $n=100$.
- Thus, the null distribution is normal with mean 0.25 and standard deviation $\sqrt{0.25*0.75/100}=0.0433$.
- The null distribution here is the distribution of correct guesses under random guessing.


Sampling Distribution Under $H_0 : p=0.25$



Difficulty of Composite Alternative Hypotheses

- Finding the sampling distribution under H_1 is impossible.
- $H_1 : p > 0.25$ only specifies a range for p .
- The true proportion p might be 0.8, it might be 0.3, it might be 0.99, we don't know.
- This is different from the “Bill vs. Maria” where the two possible models were exactly specified.

Any of the Red Curves Could Be the Alternative Distribution



Where to Place the Cutoff?

- The alternative hypothesis is designed to be the “interesting conclusion”, thus the statistical test is designed to avoid mistakenly choosing H_1 .
- Thus, we give H_0 the “benefit of the doubt” by choosing a small probability of type I error.


Why Give H_0 the Benefit of the Doubt?

- In the ESP example, we probably would not be convinced if someone did just a little better than 25%.
- While 25% is the expected number of correct guesses, you could do a little better based on chance alone.
- So how much better does the person have to do before you'd start to believe them?


Finding the Cutoff

- Again let's choose $\alpha=0.05$
- The null hypothesis contains $p=0.25$ while the alternative hypothesis contains $p>0.25$. Thus, we reject H_0 for large values of \hat{p} .
- We need 95% of the null distribution below the cutoff.

Null Distribution



Null Distribution




Blue line must
be the 95th
percentile

Finding the 95th Percentile

- The null distribution is normal with mean 0.25 and standard deviation 0.0433
- The corresponding Z-score for the 95th percentile is 1.64 and thus the cutoff is $(1.64)(0.0433)+0.25=0.3210$.
- This cutoff says:
 - Getting a little more than 25% correct out of 100 isn't enough to convince us.
 - Our benefit of the doubt says we don't believe H_1 until you get at least 32.10% correct.

Different α Values Result in Different Cutoffs



Type I and Type II Errors

	Choose H_0 (“ <u><i>do not reject</i></u> <u><i>H₀</i></u> ”)	Choose H_1 (“ <u><i>reject H₀</i></u> ”)
H_0 is true	Correct answer	WRONG! (<u>Type I error</u>)
H_1 is true	WRONG! (<u>Type II Error</u>)	Correct answer

Type I and Type II Errors

- We reject $p=0.25$ if \hat{p} is greater than 0.3210.
- The chance of making a type I error (rejecting H_0 when H_0 is in fact true) is 0.05.
- Calculating the type II error probability is more complicated.
- For each different value in the alternative, the type II error probability is different.
- For values close to 0.25 (e.g., 0.26), the type II error probability is high because the sampling distribution overlaps much with the null distribution
- For values far from 0.25 (e.g., 0.9), the probability of type II error is low.

More on Type II Error

- One issue that we will study is how many observations should we collect to achieve particular error probabilities.
- *Power and sample size calculations.*

Review

- We have n observations. The true proportion, p , is unknown.
- We have a null value of $H_0 : p=p_0$, and depending on the setting may want to test $H_1 : p>p_0$ or $H_1 : p<p_0$, or $H_1 : p\neq p_0$.
- The alternative depends on the “scientifically interesting” conclusion.
- It may only matter if p is above p_0 , or p below p_0 , or perhaps simply being different.

Review

- We use the data (specifically \hat{p} , the sample proportion) to make one of two conclusions.
- One conclusion is to reject H_0 , indicating that the data are not consistent with H_0 .
- The other possible conclusion is to “not reject H_0 ”, which indicates the data is consistent with H_0 .
- *This is not the same as saying H_0 is true, since it is impossible to distinguish, with finite data, “ $p=p_0$ ” from “ p very close to p_0 .”*

Significance Tests: Summary

- A significance test checks whether data agrees with a hypothesis
- A hypothesis is a statement about a characteristic of a variable or a collection of variables
- If the data is very unreasonable under the hypothesis, then we will reject the hypothesis
- Usually, we try to find evidence ***against*** the hypothesis

Logical Procedure

1. State a hypothesis that you would like to find evidence against
2. Get data and calculate a statistic (for example: sample mean)
3. The hypothesis (for example: population mean equals 5) determines the sampling distribution of our statistic
4. If the calculated value in 2. is very unreasonable given 3., then we conclude that the hypothesis was wrong

Significance Test

- A ***significance test*** is a way of statistically testing a hypothesis by comparing the data to values predicted by the hypothesis
- Data that fall far from the predicted values provide ***evidence against the hypothesis***

Elements of a Significance Test

- Assumptions
- Hypotheses
- Test Statistic
- P-value
- Conclusion

Assumptions

- What type of data do we have?
 - Qualitative or quantitative?
 - Different types of data require different test procedures
- What is the population distribution?
 - Is it normal? Symmetric?
 - Some tests require normal population distributions
- Which sampling method has been used?
 - We always assume simple random sampling
 - Other sampling methods are discussed in STA 675
- What is the sample size?
 - Some methods require a minimum sample size (like $n=30$)

Hypotheses

- The ***null hypothesis*** (H_0) is the hypothesis that we test (and try to find evidence against)
- The name null hypothesis refers to the fact that it often (not always) is a hypothesis of “no effect” (no effect of a medical treatment, no difference in characteristics of countries, etc.)
- The ***alternative hypothesis*** (H_a) is a hypothesis that contradicts the null hypothesis
- When we reject the null hypothesis, the alternative hypothesis is judged acceptable
- Often, the alternative hypothesis is the actual research hypothesis that we would like to “prove” by finding evidence against the null hypothesis (proof by contradiction)

Hypotheses

The hypothesis is always a statement about one or more population parameters.

Test Statistic

- The ***test statistic*** is a statistic that is calculated from the sample data
- Often, the test statistic involves a point estimator of the parameter about which the hypothesis is stated
- For example, the test statistic may involve the sample mean or sample proportion if the hypothesis is about the population mean or population proportion

P-Value

- How unusual is the observed test statistic when the null hypothesis is assumed true?
- The ***P-value*** is the probability, assuming that H_0 is true, that the test statistic takes values at least as contradictory to H_0 as the value actually observed
- The smaller the P-value, the more strongly the data contradict H_0

Conclusion

- In addition to reporting the P-value, a formal decision is made about rejecting or not rejecting the null hypothesis
- Most studies choose a cutoff of 5%.
- This corresponds to rejecting the null hypothesis for P-values smaller than 0.05.
- Smaller P-values provide more significant evidence against the null hypothesis
- “The results are significant at the 5% level”

Multiple choice

If a test of hypothesis has a Type I error probability of .05, this means that:

- a. if the null hypothesis is true, we don't reject if 5% of the time.
- b. if the null hypothesis is true, we reject it 5% of the time.
- c. if the null hypothesis is false, we don't reject it 5% of the time.
- d. if the null hypothesis is false, we reject it 5% of the time.

Significance Test for a Mean

Example

- The mean score for all high school seniors taking a college entrance exam equals 500.
- A study is conducted to see whether a different mean applies to those students born in a foreign country.
- For a random sample of 100 of such students, the mean and standard deviation on this exam equal 508 and 100.
- Q: Is mean significantly different from 500 for international students?

Significance Test for a Mean

Assumptions

- What type of data?
 - *Quantitative*
- What is the population distribution?
 - *No special assumptions.*
 - *The test refers to the population mean of the quantitative variable.*
- Which sampling method has been used?
 - *Random sampling*
- What is the sample size?
 - *Minimum sample size of $n=30$ to use Central Limit Theorem with estimated standard deviation*

Significance Test for a Mean

Hypotheses

- The null hypothesis has the form $H_0 : \mu = \mu_0$ where μ_0 is an a priori (before taking the sample) specified number like 0 or 5.3 or 500
- The most common alternative hypothesis is
$$H_1 : \mu \neq \mu_0$$
- This is called a two-sided hypothesis, since it includes values falling above and below the null hypothesis

Significance Test for a Mean

Test Statistic

- The hypothesis is about the population mean
- So, a natural test statistic would be the sample mean
- The sample mean has, for sample size of at least $n=25$, an approximately normal sampling distribution
- The parameters of the sampling distribution are, under the null hypothesis,
 - Mean = μ_0 (that is, the sampling distribution is centered around the hypothesized mean)
 - Standard error = $\frac{\sigma}{\sqrt{n}}$, estimated by $\frac{s}{\sqrt{n}}$

Significance Test for a Mean

Test Statistic

- Then, the z-score has a standard normal distribution
- The z-score measures how many estimated standard errors the sample mean falls from the hypothesized population mean
- The farther the sample mean falls from μ_0 the larger the absolute value of the z test statistic, and the stronger the evidence against the null hypothesis

$$z = \frac{\bar{Y} - \mu_0}{s / \sqrt{n}}$$

Significance Test for a Mean

p-Value

- The p -value has the advantage that different test results from different tests can be compared: The p -value is always a number between 0 and 1
- It is the probability that a standard normal distribution takes values more extreme than the observed z score
- The smaller the p -value is, the stronger is the evidence against the null hypothesis and in favor of the alternative hypothesis
- Round p -value to two or three significant digits

Example

- The mean score for all high school seniors taking a college entrance exam equals 500.
 - A study is conducted to see whether a different mean applies to those students born in a foreign country.
 - For a random sample of 100 of such students, the mean and standard deviation on this exam equal 508 and 100.
1. Set up hypotheses for a significance test.
 2. Compute the test statistic.
 3. Report the P -value, and interpret.
 4. Can you conclude that the population mean for students born in a foreign country equals 500? Why or why not?
 5. Make a decision about H_0 , using $\alpha=0.05$
 6. Construct a 95% confidence interval for μ .